

Abstract Algebra (3rd Edition)

Chapter 7.6, Problem 8E Bookmark Show all steps: on

Step-by-step solution

Step 1 of 16

Consider B be a nonempty partially ordered set, and suppose $\{A_i\}$ be a collection of abelian groups. Let us suppose that I is directed. For all $i, j \in I$, there exists $k \in I$ with $i, j \leq k$. Consider that for every pair of indices $i, j \in I$ with $i \leq j$, there is a map $\rho_{i,j}: A_i \rightarrow A_j$ such that the following hold: (1) $\rho_{i,i} = \rho_{i,i}$ whenever $i \leq j \leq k$ and (2) $\rho_{i,j} = 1$ for all $i \in I$.

Suppose $B = \bigcup A_i \times \{i\}$ be the disjoint union of the A_i . Define a relation σ on B as follows:

[Comment](#)

Step 2 of 16

$(a, i)\sigma(b, j)$ if and only if there exists $k \in I$ such that $i, j \leq k$ and $\rho_{i,k}(a) = \rho_{j,k}(b)$.

[Comment](#)

Step 3 of 16

(a) Show that σ is an equivalence relation on B . We define $\lim A_i = B/\sigma$.
 To show that σ is equivalence, we need to verify that it is reflexive, symmetric, and transitive.
 1. σ is reflexive) Consider $(a, i) \in B$. Note that $i \leq i$, and that

$$\rho_{i,i}(a) = a = \rho_{i,i}(a)$$

Thus $(a, i)\sigma(a, i)$, and hence σ is reflexive.
 2. (σ is symmetric) Consider $(a, i)\sigma(b, j)$. Then there exists $k \geq i, j$ such that $\rho_{i,k}(a) = \rho_{j,k}(b)$. Certainly $\rho_{j,k}(b) = \rho_{i,k}(a)$, so that $(b, j)\sigma(a, i)$. Hence σ is symmetric.
 3. (σ is transitive) Suppose $(a, i)\sigma(b, j)$ and $(b, j)\sigma(c, k)$. Then there exist $l \geq i, j$ such that $\rho_{i,l}(a) = \rho_{j,l}(b)$ and $m \geq j, k$ such that $\rho_{j,m}(b) = \rho_{k,m}(c)$. Since I is a directed poset, there exists $t \in I$ such that $l \geq t, m \geq t$. Now

$$\begin{aligned} \rho_{i,t}(a) &= \rho_{i,l}(\rho_{l,t}(a)) \\ &= \rho_{i,l}(\rho_{j,t}(b)) \\ &= \rho_{i,l}(b) \\ &= \rho_{i,m}(\rho_{j,m}(b)) \\ \rho_{i,t}(a) &= \rho_{i,m}(\rho_{j,m}(c)) \\ &= \rho_{i,t}(c) \end{aligned}$$

Thus $(a, i)\sigma(c, k)$, and hence σ is transitive. Hence σ is an equivalence relation.

[Comment](#)

Step 4 of 16

(b) Consider $[x]$, denote the class of x in $\lim A_i$ and define $\rho_i: A_i \rightarrow \lim A_i$ by $\rho_i(a) = [(a, i)]$. Show that if each ρ_i is injective, then ρ_i is also injective for all i . Suppose that the ρ_i are all injective. Choose $i \in I$, and consider $a, b \in A_i$ such that $\rho_i(a) = \rho_i(b)$. Then $[(a, i)] = [(b, i)]$. For some $k \geq i$, we have $\rho_{i,k}(a) = \rho_{i,k}(b)$. Since $\rho_{i,k}$ is injective, $a = b$. Hence ρ_i is injective.

[Comment](#)

Step 5 of 16

(c) Assume that the ρ_i are all group homomorphism. For $[(a, i)], [(b, j)] \in \lim A_i$, show that the operation $[(a, i)] + [(b, j)] = [(\rho_{i,k}(a) + \rho_{j,k}(b), k)]$, where k is any upper bound of i and j , is well defined and makes $\lim A_i$ an abelian group. Deduce that the ρ_i are group homomorphism.
 Consider that the ρ_i are all group homomorphism. First we show that $+$ is well defined.
 Suppose $[(a, i)] = [(a_1, i_1)]$ and $[(b, j)] = [(b_1, j_1)]$. Then there exists $s \geq i, i_1$ such that $\rho_{i,s}(a) = \rho_{i_1,s}(a_1)$ and $r \geq j, j_1$ such that $\rho_{j,r}(b) = \rho_{j_1,r}(b_1)$. Now choose arbitrary $k \geq s, r$ and $k_1 \geq s, r$. Again choose $t \geq k, k_1, r, s$. See the following:

$$\begin{aligned} \rho_{i,t}(\rho_{i,s}(a) + \rho_{j,s}(b)) &= \rho_{i,t}(\rho_{i,s}(a)) + \rho_{i,t}(\rho_{j,s}(b)) \\ &= \rho_{i,t}(\rho_{i,s}(a)) + \rho_{i,t}(\rho_{j,r}(b)) \\ &= \rho_{i,t}(\rho_{i,s}(a)) + \rho_{i,t}(\rho_{j,r}(b)) \\ \rho_{i,t}(\rho_{i,s}(a) + \rho_{j,s}(b)) &= \rho_{i,t}(\rho_{i,s}(a)) + \rho_{i,t}(\rho_{j,r}(b)) \\ &= \rho_{i,t}(\rho_{i,s}(a)) + \rho_{i,t}(\rho_{j,r}(b)) \\ &= \rho_{i,t}(\rho_{i,s}(a)) + \rho_{i,t}(\rho_{j,r}(b)) \end{aligned}$$

Hence, $(\rho_{i,t}(a) + \rho_{j,t}(b), t)\sigma(\rho_{i_1,t_1}(a_1) + \rho_{j_1,t_1}(b_1), t)$, and $[(a, i)] + [(b, j)] = [(\rho_{i,t}(a) + \rho_{j,t}(b), t)]$.

Thus $+$ is well-defined.
 Next we show that $(\lim A_i, +)$ is an abelian group.

(1) ($+$ is associative) Consider $[(a, i)], [(b, j)],$ and $[(c, k)]$ be in $\lim A_i$, and suppose $l \geq i, j, i \geq j, k$, and $m \geq l, i$. Then we have the following.

$$\begin{aligned} [(a, i)] + [(b, j)] + [(c, k)] &= [(\rho_{i,m}(a) + \rho_{j,m}(b)) + \rho_{i,m}(c), m] \\ &= [(\rho_{i,m}(a) + \rho_{j,m}(b) + \rho_{i,m}(c)), m] \\ &= [(\rho_{i,m}(a) + \rho_{i,m}(c) + \rho_{j,m}(b)), m] \\ &= [(\rho_{i,m}(a) + \rho_{i,m}(c)) + \rho_{j,m}(b), m] \\ &= [(a, i)] + [(\rho_{i,m}(a) + \rho_{i,m}(c)), m] \\ &= [(a, i)] + [((b, j)) + [(c, k)]] \end{aligned}$$

So $+$ is associative.

[Comment](#)

Step 6 of 16

(2) For all $i, j \in I$, there exists $k \geq i, j$, and $\rho_{i,k}(0) = \rho_{j,k}(0)$ since the ρ_i are group homomorphism. Hence $[(0, i)] = [(0, j)]$ for all i, j . Consider $0 = [(0, i)]$. Suppose $[(a, i)] \in \lim A_i$. Then

$$\begin{aligned} 0 + [(a, i)] &= [(0, i)] + [(a, i)] \\ &= [(\rho_{i,i}(0) + \rho_{i,i}(a)), i] \\ &= [(0 + a, i)] \\ &= [(a, i)] \end{aligned}$$

Thus, $[(a, i)] + 0 = [(a, i)]$. Hence $0 = [(0, i)]$ is an additive identity element.

[Comment](#)

Step 7 of 16

3. Consider $[(a, i)] \in \lim A_i$. Note that $[(a, i)] + [(-a, i)] = [(\rho_{i,i}(a) + \rho_{i,i}(-a)), i] = [(a - a, i)] = [(0, i)] = 0$. Hence every element of $\lim A_i$ has an additive inverse.

[Comment](#)

Step 8 of 16

4. Suppose $[(a, i)], [(b, j)] \in \lim A_i$, and consider $k \geq i, j$. Then $[(a, i)] + [(b, j)] = [(\rho_{i,k}(a) + \rho_{j,k}(b), k)] = [(\rho_{i,k}(a) + \rho_{j,k}(b), k)] = [(a, i)] + [(b, j)]$. Hence $+$ is commutative.

Thus $(\lim A_i, +)$ is an abelian group. Finally, we show that each $\rho_i: A_i \rightarrow \lim A_i$ is a group homomorphism. Suppose $a, b \in A_i$. Then

$$\begin{aligned} \rho_i(a + b) &= [(a + b, i)] \\ &= [(\rho_{i,i}(a) + \rho_{i,i}(b)), i] \\ &= [(a, i)] + [(b, i)] \\ &= \rho_i(a) + \rho_i(b) \end{aligned}$$

Hence ρ_i is a group homomorphism for all i .

[Comment](#)

Step 9 of 16

(d) Prove that if all the A_i are commutative rings with $1 \neq 0$ and all the ρ_i are unital ring homomorphism, then $\lim A_i$ may likewise be given the structure of a commutative ring with $1 \neq 0$ such that the ρ_i are all ring homomorphism.
 Define an operator on $\lim A_i$ as follows: $[(a, i)] \cdot [(b, j)] = [(\rho_{i,k}(a) \cdot \rho_{j,k}(b), k)]$, where k is any upper bound of i and j in I . See this.

(1) (\cdot is well defined) Consider $[(a, i)] = [(a_1, i_1)]$ and $[(b, j)] = [(b_1, j_1)]$. Then there exist $r \geq i, i_1$ and $s \geq j, j_1$ such that $\rho_{i,r}(a) = \rho_{i_1,r}(a_1)$ and $\rho_{j,s}(b) = \rho_{j_1,s}(b_1)$. Choose $k \geq r, s$ and $k_1 \geq r, s$, and $t \geq k, k_1$. Now

$$\begin{aligned} \rho_{i,t}(\rho_{i,r}(a) \cdot \rho_{j,r}(b)) &= \rho_{i,t}(\rho_{i,r}(a)) \cdot \rho_{i,t}(\rho_{j,r}(b)) \\ &= \rho_{i,t}(\rho_{i,r}(a)) \cdot \rho_{i,t}(\rho_{j,s}(b)) \\ &= \rho_{i,t}(\rho_{i,r}(a)) \cdot \rho_{i,t}(\rho_{j,s}(b)) \\ \rho_{i,t}(\rho_{i,r}(a) \cdot \rho_{j,r}(b)) &= \rho_{i,t}(\rho_{i,r}(a)) \cdot \rho_{i,t}(\rho_{j,s}(b)) \\ &= \rho_{i,t}(\rho_{i,r}(a)) \cdot \rho_{i,t}(\rho_{j,s}(b)) \end{aligned}$$

Thus $(\rho_{i,t}(a) \cdot \rho_{j,t}(b), t)\sigma(\rho_{i_1,t_1}(a_1) \cdot \rho_{j_1,t_1}(b_1), t)$, and particularly, $[(a, i)] \cdot [(b, j)] = [(a_1, i_1)] \cdot [(b_1, j_1)]$. So \cdot is well-defined.

[Comment](#)

Step 10 of 16

(2) (\cdot is associative) Suppose $[(a, i)], [(b, j)], [(c, k)] \in \lim A_i$. Consider $r \geq i, j, i \geq j, k$, and $t \geq r, s$. Then we have the following:

$$\begin{aligned} [(a, i)] \cdot [(b, j)] \cdot [(c, k)] &= [(\rho_{i,t}(a) \cdot \rho_{j,t}(b)) \cdot \rho_{i,t}(c), t] \\ &= [(\rho_{i,t}(a) \cdot \rho_{j,t}(b) \cdot \rho_{i,t}(c)), t] \\ &= [(\rho_{i,t}(a) \cdot \rho_{i,t}(c) \cdot \rho_{j,t}(b)), t] \\ &= [(\rho_{i,t}(a) \cdot \rho_{i,t}(c)) \cdot \rho_{j,t}(b), t] \\ &= [(a, i)] \cdot [(\rho_{i,t}(a) \cdot \rho_{i,t}(c)), t] \\ &= [(a, i)] \cdot [((b, j)) \cdot [(c, k)]] \end{aligned}$$

Hence \cdot is associative.

[Comment](#)

Step 11 of 16

(3) (\cdot distributes over $+$) We will show that \cdot distributes over $+$ on the left; distributes on the right is similar. Let $[(a, i)], [(b, j)], [(c, k)] \in \lim A_i$. Suppose $r \geq j, k$, and let $t \geq i, r$. Then we have the following:

$$\begin{aligned} [(a, i)] \cdot [((b, j)) + [(c, k)]] &= [(a, i)] \cdot [(\rho_{j,t}(b) + \rho_{k,t}(c), t)] \\ &= [(\rho_{i,t}(a) \cdot \rho_{j,t}(b) + \rho_{i,t}(a) \cdot \rho_{k,t}(c)), t] \\ &= [(\rho_{i,t}(a) \cdot \rho_{j,t}(b) + \rho_{i,t}(a) \cdot \rho_{k,t}(c)), t] \\ &= [(\rho_{i,t}(a) \cdot \rho_{j,t}(b) + \rho_{i,t}(a) \cdot \rho_{k,t}(c)), t] \\ &= [(\rho_{i,t}(a) \cdot \rho_{j,t}(b) + \rho_{i,t}(a) \cdot \rho_{k,t}(c)), t] \\ &= [(a, i)] \cdot [((b, j)) + [(c, k)]] \end{aligned}$$

Hence \cdot distributes over $+$.

Thus $(\lim A_i, +, \cdot)$ is a ring.

[Comment](#)

Step 12 of 16

However, we have the following:
 (1) Consider the A_i are all commutative. Suppose $[(a, i)], [(b, j)] \in \lim A_i$, and let $k \geq i, j$. Then we have the following:

$$\begin{aligned} [(a, i)] \cdot [(b, j)] &= [(\rho_{i,k}(a) \cdot \rho_{j,k}(b), k)] \\ &= [(\rho_{i,k}(b) \cdot \rho_{j,k}(a), k)] \\ &= [(b, j)] \cdot [(a, i)] \end{aligned}$$

Hence $\lim A_i$ is a commutative ring. If all the A_i are commutative, then $\lim A_i$ is commutative.
 (2) Note that because the ρ_i are unital ring homomorphism

$$\rho_{i,i}(1) = 1 = \rho_{i,i}(1)$$

Whenever $k \geq i, j$. Hence $[(1, i)] = [(1, j)]$ for all i, j . Define $1 = [(1, i)]$. Consider $[(a, i)] \in \lim A_i$. Then,

$$\begin{aligned} 1 \cdot [(a, i)] &= [(1, i)] \cdot [(a, i)] \\ &= [(\rho_{i,i}(1) \cdot \rho_{i,i}(a)), i] \\ &= [(1, i)] \cdot [(a, i)] \\ &= [(a, i)] \end{aligned}$$

Thus, $[(a, i)] \cdot 1 = [(a, i)]$. Hence 1 is a multiplicative identity in $\lim A_i$. If all the A_i have $1 \neq 0$ and the ρ_i are unital, then $\lim A_i$ has a multiplicative identity.
 3. Suppose $[(0, i)] = [(0, j)]$. Then there exists $j \geq i$ such that $\rho_{i,j}(0) = \rho_{j,i}(0)$, so that $0 = 1$ in A_i a contradiction. Hence $1 \neq 0$ in $\lim A_i$.
 Thus if the A_i are commutative rings with $1 \neq 0$, then $\lim A_i$ is a commutative ring with $1 \neq 0$. Thus, if $1 \neq 0$ for all A_i , then $1 \neq 0$ in $\lim A_i$.

[Comment](#)

Step 13 of 16

Therefore, that if all the A_i are commutative rings with $1 \neq 0$ and all the ρ_i are unital ring homomorphism, then $\lim A_i$ may likewise be given the structure of a commutative ring with $1 \neq 0$ such that the ρ_i are all ring homomorphism.

[Comment](#)

Step 14 of 16

(e) Under the hypotheses of part (c), prove that $\lim A_i$ has the following universal property: if C is any abelian group such that for each $i \in I$ there is a homomorphism $\varphi_i: A_i \rightarrow C$ with $\varphi_i \circ \rho_{i,j} = \varphi_j$ whenever $i \leq j$, then there is a unique homomorphism $\varphi: \lim A_i \rightarrow C$ such that $\varphi \circ \rho_i = \varphi_i$ for all i .
 Consider, C is an abelian group and that we have an indexed family of group homomorphism $\varphi_i: A_i \rightarrow C$ such that $\varphi_i \circ \rho_{i,j} = \varphi_j$ for all $i, j \in I$.
 Define $\varphi: \lim A_i \rightarrow C$ by $\varphi([(a, i)]) = \varphi_i(a)$. We need to show that φ is a well defined group homomorphism.
 Suppose $[(a, i)] = [(b, j)]$. Then there exists $k \geq i, j$ such that $\rho_{i,k}(a) = \rho_{j,k}(b)$. Since $\rho_{i,k}(a) = \rho_{i,k}(b)$ and we have $\varphi_i(\rho_{i,k}(a)) = \varphi_i(\rho_{i,k}(b))$. Then, $(\varphi_i \circ \rho_{i,k})(a) = (\varphi_i \circ \rho_{i,k})(b)$, and we have $\varphi_i(a) = \varphi_i(b)$. Hence, $\varphi([(a, i)]) = \varphi([(b, j)])$, and φ is well defined.
 Hence, $\varphi \circ \rho_i = \varphi_i$ for all i .

[Comment](#)

Step 16 of 16

Suppose that we have a group homomorphism $\psi: \lim A_i \rightarrow C$ which also satisfies $\psi \circ \rho_i = \varphi_i$ for all $i \in I$. Then for all i and all $a \in A_i$.

$$\begin{aligned} \psi([(a, i)]) &= \psi(\rho_i(a)) \\ &= (\psi \circ \rho_i)(a) \\ &= \varphi_i(a) \\ &= (\varphi \circ \rho_i)(a) \\ \psi([(a, i)]) &= \varphi(\rho_i(a)) \\ &= \varphi([(a, i)]) \end{aligned}$$

Hence $\psi = \varphi$.
 Therefore, $\lim A_i$ has the following universal property: if C is any abelian group such that for each $i \in I$ there is a homomorphism $\varphi_i: A_i \rightarrow C$ with $\varphi_i \circ \rho_{i,j} = \varphi_j$ whenever $i \leq j$, then there is a unique homomorphism $\varphi: \lim A_i \rightarrow C$ such that $\varphi \circ \rho_i = \varphi_i$ for all i .

[Comment](#)

Was this solution helpful? 1 0

Post a question
 Answers from our experts for your tough homework questions.

Continue to edit and attach image(s).

My Textbook Solutions

Abstract Algebra 3rd Edition	Computer Security 3rd Edition	Classical Mechanics 3rd Edition

[View all solutions](#)

Chegg tutors who can help right now

	Ileana University of Virginia	341
	Rahul Chartered Account...	178
	Aman JAMIA MILLIA ISLIA...	215

- | | | | | | |
|--|---|-----------------------------------|--------------------------------------|---------------------------------|--|
| Become a Tutor | Advertising Choices | Cheap Textbooks | Online Tutoring | EasyBib | Customer Service |
| Chegg For Good | Cookie Notice | Chegg Coupon | Sell Textbooks | Internships.com | Give Us Feedback |
| College Marketing | General Policies | Chegg Play | Solutions Manual | StudyBlue | Help with Chegg Tutors |
| Corporate Development | Intellectual Property Rights | Chegg Study Help | Study 101 | | Help with eBooks |
| Investor Relations | International Privacy Policy | College Textbooks | Test Prep | | Help to use EasyBib Plus |
| Jobs | Terms of Use | eTextbooks | Textbook Rental | | Marissa Chegg Study Subscription |
| Join Our Affiliate Program | Chegg Tutors Terms of Service | Chegg Math Solver | Used Textbooks | | Return Your Books |
| Media Center | US Privacy Policy | Mobile Apps | Digital Access Codes | | Textbook Return Policy |
| Site Map | Your CA Privacy Rights | | | | |
| | Honor Code | | | | |



© 2003-2019 Chegg Inc. All rights reserved.