

Table of Contents

Review Questions Chapter 1	2
Solution Manual Chapter 1	4
1 Questions	4
2 Exercises	6
2.1 Development of the Atomic Theory	6
2.2 The Nature of the Atom	8
2.3 Chem Work Problems	12
3 Challenge Problems	15
3.1 Marathon Problem	17

Review Questions – Chapter 1

1.
 - a. Law versus theory: A law is a concise statement or equation that summarizes observed behavior. A theory is a set of hypotheses that gives an overall explanation of some phenomenon. A law summarizes what happens; a theory (or model) attempts to explain why it happens.
 - b. Theory versus experiment: A theory is an explanation of why things behave the way they do, while an experiment is the process of observing that behavior. Theories attempt to explain the results of experiments and are, in turn, tested by further experiments.
 - c. Qualitative versus quantitative: A qualitative observation only describes a quality while a quantitative observation attaches a number to the observation. Some qualitative observations would be: The water was hot to the touch. Mercury was found in the drinking water. Some quantitative observations would be: The temperature of the water was 62°C. The concentration of mercury in the drinking water was 1.5 ppm.
 - d. Hypothesis versus theory: Both are explanations of experimental observation. A theory is a set of hypotheses that has been tested over time and found to still be valid, with (perhaps) some modifications.
2. No, it is useful whenever a systematic approach of observation and hypothesis testing can be used.
3.
 - a. Atoms have specific masses and are neither created nor destroyed by chemical reactions. Because atoms are conserved in a chemical reaction, mass cannot change in a chemical reaction. Mass is conserved.
 - b. The composition of a substance depends on the number and kinds of atoms that form it. A certain compound always has the same number and kinds of atoms in its formula.
 - c. Compounds of the same elements differ only in the numbers of atoms of the elements forming them, i.e., NO, N₂O, NO₂.
4. Deflection of cathode rays by magnetic and electric fields led to the conclusion that cathode rays were negatively charged. The cathode ray was produced at the negative electrode and repelled by the negative pole of the applied electric field.
5. J. J. Thomson discovered electrons. He postulated that all atoms must contain electrons, but Thomson also postulated that atoms must contain positive charge in order for the atom to be electrically neutral. Henri Becquerel discovered radioactivity. Lord Rutherford proposed the nuclear model of the atom. Dalton's original model proposed that atoms were indivisible particles (that is, atoms had no internal structure). Thomson and Becquerel discovered subatomic particles, and Rutherford's model attempted to describe the internal structure of the atom composed of these subatomic particles. In addition, the existence of isotopes, atoms of the same element but with different mass, had to be included in the model.
6. If the plum pudding model were correct (a diffuse positive charge with electrons scattered throughout), then alpha particles should have traveled through the thin foil with very minor

deflections in their path. This was not the case as a few of the alpha particles were deflected at very large angles. Rutherford reasoned that the large deflections of these alpha particles could be caused only by a center of concentrated positive charge that contains most of the atom's mass (the nuclear model of the atom).

7. The proton and neutron have similar mass with the mass of the neutron slightly larger than that of the proton. Each of these particles has a mass approximately 1800 times greater than that of an electron. The combination of the protons and the neutrons in the nucleus makes up the bulk of the mass of an atom, but the electrons make the greatest contribution to the chemical properties of the atom.
8. The atomic number of an element is equal to the number of protons in the nucleus of an atom of that element. The mass number is the sum of the number of protons plus neutrons in the nucleus. The atomic mass is the actual mass of a particular isotope (including electrons). As we will see in Chapter 5, the average mass of an atom is taken from a measurement made on a large number of atoms. The average atomic mass value is listed in the periodic table.

CHAPTER 1

CHEMICAL FOUNDATIONS

Questions

11. A law summarizes what happens, e.g., law of conservation of mass in a chemical reaction or the ideal gas law, $PV = nRT$. A theory (model) is an attempt to explain why something happens. Dalton's atomic theory explains why mass is conserved in a chemical reaction. The kinetic molecular theory explains why pressure and volume are inversely related at constant temperature and moles of gas present, as well as explaining the other mathematical relationships summarized in $PV = nRT$.
12. a. At 8 a.m., approximately 57 cars pass through the intersection per hour.
b. At 12 a.m. (midnight), only 1 or 2 cars pass through the intersection per hour.
c. Traffic at the intersection is limited to less than 10 cars per hour from 8 p.m. to 5 a.m. Starting at 6 a.m., there is a steady increase in traffic through the intersection, peaking at 8 a.m. when approximately 57 cars pass per hour. Past 8 a.m. traffic moderates to about 40 cars through the intersection per hour until noon, and then decreases to 21 cars per hour by 3 p.m. Past 3 p.m. traffic steadily increases to a peak of 52 cars per hour at 5 p.m., and then steadily decreases to the overnight level of less than 10 cars through the intersection per hour.
d. The traffic pattern through the intersection is directly related to the work schedules of the general population as well as to the store hours of the businesses in downtown.
e. Run the same experiment on a Sunday, when most of the general population doesn't work and when a significant number of downtown stores are closed in the morning.
13. The fundamental steps are
(1) making observations;
(2) formulating hypotheses;
(3) performing experiments to test the hypotheses.

The key to the scientific method is performing experiments to test hypotheses. If after the test of time the hypotheses seem to account satisfactorily for some aspect of natural behavior, then the set of tested hypotheses turns into a theory (model). However, scientists continue to perform experiments to refine or replace existing theories. Hence, science is a dynamic or active process, not a static one.

14. A compound will always contain the same numbers (and types) of atoms. A given amount of hydrogen will react only with a specific amount of oxygen. Any excess oxygen will remain unreacted.
15. Law of conservation of mass: Mass is neither created nor destroyed. The total mass before a chemical reaction always equals the total mass after a chemical reaction.

CHAPTER 1 CHEMICAL FOUNDATIONS

Law of definite proportion: A given compound always contains exactly the same proportion of elements by mass. For example, water is always 1 g H for every 8 g oxygen.

Law of multiple proportions: When two elements form a series of compounds, the ratios of the mass of the second element that combine with 1 g of the first element always can be reduced to small whole numbers. For CO_2 and CO discussed in Section 1.4, the mass ratios of oxygen that react with 1 g carbon in each compound are in a 2 : 1 ratio.

16. Yes, 1.0 g H would react with 37.0 g ^{37}Cl , and 1.0 g H would react with 35.0 g ^{35}Cl .
- No, the mass ratio of H/Cl would always be 1 g H/37 g Cl for ^{37}Cl and 1 g H/35 g Cl for ^{35}Cl . As long as we had pure ^{37}Cl or pure ^{35}Cl , the ratios will always hold. If we have a mixture (such as the natural abundance of chlorine), the ratio will also be constant as long as the composition of the mixture of the two isotopes does not change.
17. Natural niacin and commercially produced niacin have the exact same formula of $\text{C}_6\text{H}_5\text{NO}_2$. Therefore, both sources produce niacin having an identical nutritional value. There may be other compounds present in natural niacin that would increase the nutritional value, but the nutritional value due to just niacin is identical to the commercially produced niacin.
18. a. The smaller parts are electrons and the nucleus. The nucleus is broken down into protons and neutrons, which can be broken down into quarks. For our purpose, electrons, neutrons, and protons are the key smaller parts of an atom.
- b. All atoms of hydrogen have 1 proton in the nucleus. Different isotopes of hydrogen have 0, 1, or 2 neutrons in the nucleus. Because we are talking about atoms, this implies a neutral charge, which dictates 1 electron present for all hydrogen atoms. If charged ions were included, then different ions/atoms of H could have different numbers of electrons.
- c. Hydrogen atoms always have 1 proton in the nucleus, and helium atoms always have 2 protons in the nucleus. The number of neutrons can be the same for a hydrogen atom and a helium atom. Tritium (^3H) and ^4He both have 2 neutrons. Assuming neutral atoms, then the number of electrons will be 1 for hydrogen and 2 for helium.
- d. Water (H_2O) is always 1 g hydrogen for every 8 g of O present, whereas H_2O_2 is always 1 g hydrogen for every 16 g of O present. These are distinctly different compounds, each with its own unique relative number and types of atoms present.
- e. A chemical equation involves a reorganization of the atoms. Bonds are broken between atoms in the reactants, and new bonds are formed in the products. The number and types of atoms between reactants and products do not change. Because atoms are conserved in a chemical reaction, mass is also conserved.
19. J. J. Thomson's study of cathode-ray tubes led him to postulate the existence of negatively charged particles that we now call electrons. Ernest Rutherford and his alpha bombardment of metal foil experiments led him to postulate the nuclear atom—an atom with a tiny dense center of positive charge (the nucleus) with electrons moving about the nucleus at relatively large distances away; the distance is so large that an atom is mostly empty space.


CHAPTER 1 CHEMICAL FOUNDATIONS

20. The atom is composed of a tiny dense nucleus containing most of the mass of the atom. The nucleus itself is composed of neutrons and protons. Neutrons have a mass slightly larger than that of a proton and have no charge. Protons, on the other hand, have a 1+ relative charge as compared to the 1- charged electrons; the electrons move about the nucleus at relatively large distances. The volume of space that the electrons move about is so large, as compared to the nucleus, that we say an atom is mostly empty space.
21. The number and arrangement of electrons in an atom determine how the atom will react with other atoms. The electrons determine the chemical properties of an atom. The number of neutrons present determines the isotope identity.
22. Density = mass/volume; if the volumes are assumed equal, then the much more massive proton would have a much larger density than the relatively light electron.
23. For lighter, stable isotopes, the number of protons in the nucleus is about equal to the number of neutrons. When the number of protons and neutrons is equal to each other, the mass number (protons + neutrons) will be twice the atomic number (protons). Therefore, for lighter isotopes, the ratio of the mass number to the atomic number is close to 2. For example, consider ^{28}Si , which has 14 protons and $(28 - 14 =) 14$ neutrons. Here, the mass number to atomic number ratio is $28/14 = 2.0$. For heavier isotopes, there are more neutrons than protons in the nucleus. Therefore, the ratio of the mass number to the atomic number increases steadily upward from 2 as the isotopes get heavier and heavier. For example, ^{238}U has 92 protons and $(238 - 92 =) 146$ neutrons. The ratio of the mass number to the atomic number for ^{238}U is $238/92 = 2.6$.
24. Some elements exist as molecular substances. That is, hydrogen normally exists as H_2 molecules, not single hydrogen atoms. The same is true for N_2 , O_2 , F_2 , Cl_2 , Br_2 , and I_2 .

Exercises

Development of the Atomic Theory

25.
 - a. The composition of a substance depends on the numbers of atoms of each element making up the compound (depends on the formula of the compound) and not on the composition of the mixture from which it was formed.
 - b. Avogadro's hypothesis (law) implies that volume ratios are equal to molecule ratios at constant temperature and pressure. $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow 2 \text{HCl}(\text{g})$. From the balanced equation, the volume of HCl produced will be twice the volume of H_2 (or Cl_2) reacted.
26. Avogadro's hypothesis (law) implies that volume ratios are equal to molecule ratios at constant temperature and pressure. Here, 1 volume of N_2 reacts with 3 volumes of H_2 to produce 2 volumes of the gaseous product or in terms of molecule ratios:



In order for the equation to be balanced, the product must be NH_3 .

CHAPTER 1 CHEMICAL FOUNDATIONS

27. From the law of definite proportions, a given compound always contains exactly the same proportion of elements by mass. The first sample of chloroform has a total mass of 12.0 g C + 106.4 g Cl + 1.01 g H = 119.41 g (carrying extra significant figures). The mass percent of carbon in this sample of chloroform is:

$$\frac{12.0 \text{ g C}}{119.41 \text{ g total}} \times 100 = 10.05\% \text{ C by mass}$$

From the law of definite proportions, the second sample of chloroform must also contain 10.05% C by mass. Let x = mass of chloroform in the second sample:

$$\frac{30.0 \text{ g C}}{x} \times 100 = 10.05, x = 299 \text{ g chloroform}$$


28. A compound will always have a constant composition by mass. From the initial data given, the mass ratio of H : S : O in sulfuric acid (H_2SO_4) is:

$$\frac{2.02}{2.02} : \frac{32.07}{2.02} : \frac{64.00}{2.02} = 1 : 15.9 : 31.7$$

If we have 7.27 g H, then we will have $7.27 \times 15.9 = 116 \text{ g S}$ and $7.27 \times 31.7 = 230. \text{ g O}$ in the second sample of H_2SO_4 .

29. Mass is conserved in a chemical reaction because atoms are conserved. Chemical reactions involve the reorganization of atoms, so formulas change in a chemical reaction, but the number and types of atoms do not change. Because the atoms do not change in a chemical reaction, mass must not change. In this equation we have two oxygen atoms and four hydrogen atoms both before and after the reaction occurs.

30. Mass is conserved in a chemical reaction.



Mass of reactants = $46.0 + 96.0 = 142.0 \text{ g}$ = mass of products

$$142.0 \text{ g} = 54.0 \text{ g} + \text{mass of CO}_2, \text{ mass of CO}_2 = 142.0 - 54.0 = 88.0 \text{ g}$$

31. To get the atomic mass of H to be 1.00, we divide the mass of hydrogen that reacts with 1.00 g of oxygen by 0.126; that is, $\frac{0.126}{0.126} = 1.00$. To get Na, Mg, and O on the same scale, we do the same division.

$$\text{Na: } \frac{2.875}{0.126} = 22.8; \text{ Mg: } \frac{1.500}{0.126} = 11.9; \text{ O: } \frac{1.00}{0.126} = 7.94$$

	H	O	Na	Mg
Relative value	1.00	7.94	22.8	11.9
Accepted value	1.008	16.00	22.99	24.31

For your information, the atomic masses of O and Mg are incorrect. The atomic masses of H and Na are close to the values given in the periodic table. Something must be wrong about the assumed formulas of the compounds. It turns out the correct formulas are H_2O , Na_2O , and MgO . The smaller discrepancies result from the error in the assumed atomic mass of H.

CHAPTER 1 CHEMICAL FOUNDATIONS

32. If the formula is InO , then one atomic mass of In would combine with one atomic mass of O, or:

$$\frac{A}{16.00} = \frac{4.784 \text{ g In}}{1.000 \text{ g O}}, A = \text{atomic mass of In} = 76.54$$

If the formula is In_2O_3 , then two times the atomic mass of In will combine with three times the atomic mass of O, or:

$$\frac{2A}{(3)16.00} = \frac{4.784 \text{ g In}}{1.000 \text{ g O}}, A = \text{atomic mass of In} = 114.8$$

The latter number is the atomic mass of In used in the modern periodic table.

The Nature of the Atom

33. From section 1-7, the nucleus has “a diameter of about 10^{-13} cm ” and the electrons “move about the nucleus at an average distance of about 10^{-8} cm from it.” We will use these statements to help determine the densities. Density of hydrogen nucleus (contains one proton only):

$$V_{\text{nucleus}} = \frac{4}{3} \pi r^3 = \frac{4}{3} (3.14) (5 \times 10^{-14} \text{ cm})^3 = 5 \times 10^{-40} \text{ cm}^3$$

$$d = \text{density} = \frac{1.67 \times 10^{-24} \text{ g}}{5 \times 10^{-40} \text{ cm}^3} = 3 \times 10^{15} \text{ g/cm}^3$$

Density of H atom (contains one proton and one electron):

$$V_{\text{atom}} = \frac{4}{3} (3.14) (1 \times 10^{-8} \text{ cm})^3 = 4 \times 10^{-24} \text{ cm}^3$$

$$d = \frac{1.67 \times 10^{-24} \text{ g} + 9 \times 10^{-28} \text{ g}}{4 \times 10^{-24} \text{ cm}^3} = 0.4 \text{ g/cm}^3$$

34. Because electrons move about the nucleus at an average distance of about $1 \times 10^{-8} \text{ cm}$, the diameter of an atom will be about $2 \times 10^{-8} \text{ cm}$. Let's set up a ratio:

$$\frac{\text{diameter of nucleus}}{\text{diameter of atom}} = \frac{1 \text{ mm}}{\text{diameter of model}} = \frac{1 \times 10^{-13} \text{ cm}}{2 \times 10^{-8} \text{ cm}}; \text{ solving:}$$

$$\text{diameter of model} = 2 \times 10^5 \text{ mm} = 200 \text{ m}$$

35. $3 \times 5.93 \times 10^{-18} \text{ C} \times \frac{1 \text{ electron charge}}{1.602 \times 10^{-19} \text{ C}} = 37 \text{ negative (electron) charges on the oil drop}$

36. First, divide all charges by the smallest quantity, 6.40×10^{-13} :

$$\frac{2.56 \times 10^{-12}}{6.40 \times 10^{-13}} = 4.00; \quad \frac{7.68}{0.640} = 12.0; \quad \frac{3.84}{0.640} = 6.00$$

CHAPTER 1 CHEMICAL FOUNDATIONS

Because all charges are whole-number multiples of 6.40×10^{-13} zirkombs, the charge on one electron could be 6.40×10^{-13} zirkombs. However, 6.40×10^{-13} zirkombs could be the charge of two electrons (or three electrons, etc.). All one can conclude is that the charge of an electron is 6.40×10^{-13} zirkombs or an integer fraction of 6.40×10^{-13} zirkombs.

37. Z is the atomic number and is equal to the number of protons in the nucleus. A is the mass number and is equal to the number of protons plus neutrons in the nucleus. X is the symbol of the element. See the front cover of the text which has a listing of the symbols for the various elements and corresponding atomic number or see the periodic table on the cover to determine the identity of the various atoms. Because all of the atoms have equal numbers of protons and electrons, each atom is neutral in charge.
- a. $^{23}_{11}\text{Na}$ b. $^{19}_9\text{F}$ c. $^{16}_8\text{O}$
38. The atomic number for carbon is 6. ^{14}C has 6 protons, $14 - 6 = 8$ neutrons, and 6 electrons in the neutral atom. ^{12}C has 6 protons, $12 - 6 = 6$ neutrons, and 6 electrons in the neutral atom. The only difference between an atom of ^{14}C and an atom of ^{12}C is that ^{14}C has two additional neutrons.
39. a. $^{79}_{35}\text{Br}$: 35 protons, $79 - 35 = 44$ neutrons. Because the charge of the atom is neutral, the number of protons = the number of electrons = 35.
- b. $^{81}_{35}\text{Br}$: 35 protons, 46 neutrons, 35 electrons
- c. $^{239}_{94}\text{Pu}$: 94 protons, 145 neutrons, 94 electrons
- d. $^{133}_{55}\text{Cs}$: 55 protons, 78 neutrons, 55 electrons
- e. ^3_1H : 1 proton, 2 neutrons, 1 electron
- f. $^{56}_{26}\text{Fe}$: 26 protons, 30 neutrons, 26 electrons
40. a. $^{235}_{92}\text{U}$: 92 p, 143 n, 92 e b. $^{27}_{13}\text{Al}$: 13 p, 14 n, 13 e c. $^{57}_{26}\text{Fe}$: 26 p, 31 n, 26 e
- d. $^{208}_{82}\text{Pb}$: 82 p, 126 n, 82 e e. $^{86}_{37}\text{Rb}$: 37 p, 49 n, 37 e f. $^{41}_{20}\text{Ca}$: 20 p, 21 n, 20 e
41. a. Element 8 is oxygen. $A = \text{mass number} = 9 + 8 = 17$; $^{17}_8\text{O}$
- b. Chlorine is element 17. $^{37}_{17}\text{Cl}$ c. Cobalt is element 27. $^{60}_{27}\text{Co}$
- d. $Z = 26$; $A = 26 + 31 = 57$; $^{57}_{26}\text{Fe}$ e. Iodine is element 53. $^{131}_{53}\text{I}$
- f. Lithium is element 3. ^7_3Li

CHAPTER 1 CHEMICAL FOUNDATIONS

42. a. Cobalt is element 27. $A = \text{mass number} = 27 + 31 = 58$; ${}^{58}_{27}\text{Co}$
- b. ${}^{10}_5\text{B}$ c. ${}^{23}_{12}\text{Mg}$ d. ${}^{132}_{53}\text{I}$ e. ${}^{47}_{20}\text{Ca}$ f. ${}^{65}_{29}\text{Cu}$
43. a. Ba is element 56. Ba^{2+} has 56 protons, so Ba^{2+} must have 54 electrons in order to have a net charge of $2+$.
- b. Zn is element 30. Zn^{2+} has 30 protons and 28 electrons.
- c. N is element 7. N^{3-} has 7 protons and 10 electrons.
- d. Rb is element 37, Rb^{+} has 37 protons and 36 electrons.
- e. Co is element 27. Co^{3+} has 27 protons and 24 electrons.
- f. Te is element 52. Te^{2-} has 52 protons and 54 electrons.
- g. Br is element 35. Br^{-} has 35 protons and 36 electrons.
44. a. ${}^{24}_{12}\text{Mg}$: 12 protons, 12 neutrons, 12 electrons
- b. ${}^{24}_{12}\text{Mg}^{2+}$: 12 p, 12 n, 10 e c. ${}^{59}_{27}\text{Co}^{2+}$: 27 p, 32 n, 25 e
- d. ${}^{59}_{27}\text{Co}^{3+}$: 27 p, 32 n, 24 e e. ${}^{59}_{27}\text{Co}$: 27 p, 32 n, 27 e
- f. ${}^{79}_{34}\text{Se}$: 34 p, 45 n, 34 e g. ${}^{79}_{34}\text{Se}^{2-}$: 34 p, 45 n, 36 e
- h. ${}^{63}_{28}\text{Ni}$: 28 p, 35 n, 28 e i. ${}^{59}_{28}\text{Ni}^{2+}$: 28 p, 31 n, 26 e
45. Atomic number = 63 (Eu); net charge = $+63 - 60 = 3+$; mass number = $63 + 88 = 151$; symbol: ${}^{151}_{63}\text{Eu}^{3+}$
- Atomic number = 50 (Sn); mass number = $50 + 68 = 118$; net charge = $+50 - 48 = 2+$;
 symbol: ${}^{118}_{50}\text{Sn}^{2+}$
46. Atomic number = 16 (S); net charge = $+16 - 18 = 2-$; mass number = $16 + 18 = 34$; symbol: ${}^{34}_{16}\text{S}^{2-}$
- Atomic number = 16 (S); net charge = $+16 - 18 = 2-$; mass number = $16 + 16 = 32$;
 symbol: ${}^{32}_{16}\text{S}^{2-}$

CHAPTER 1 CHEMICAL FOUNDATIONS

47.

Symbol	Number of protons in nucleus	Number of neutrons in nucleus	Number of electrons	Net charge
${}_{92}^{238}\text{U}$	92	146	92	0
${}_{20}^{40}\text{Ca}^{2+}$	20	20	18	2+
${}_{23}^{51}\text{V}^{3+}$	23	28	20	3+
${}_{39}^{89}\text{Y}$	39	50	39	0
${}_{35}^{79}\text{Br}^{-}$	35	44	36	1-
${}_{15}^{31}\text{P}^{3-}$	15	16	18	3-

48.

Symbol	Number of protons in nucleus	Number of neutrons in nucleus	Number of electrons	Net charge
${}_{26}^{53}\text{Fe}^{2+}$	26	27	24	2+
${}_{26}^{59}\text{Fe}^{3+}$	26	33	23	3+
${}_{85}^{210}\text{At}^{-}$	85	125	86	1-
${}_{13}^{27}\text{Al}^{3+}$	13	14	10	3+
${}_{52}^{128}\text{Te}^{2-}$	52	76	54	2-

CHAPTER 1 CHEMICAL FOUNDATIONS

Chem Work Problems

49.

Number of protons in nucleus	Number of neutrons in nucleus	Symbol
9	10	$^{19}_{9}\text{F}$
13	14	$^{27}_{13}\text{Al}$
53	74	$^{127}_{53}\text{I}$
34	45	$^{79}_{34}\text{Se}$
16	16	$^{32}_{16}\text{S}$

50.

Symbol	Number of protons in nucleus	Number of neutrons in nucleus
^4_2He	2	2
$^{20}_{10}\text{Ne}$	10	10
$^{48}_{22}\text{Ti}$	22	26
$^{190}_{76}\text{Os}$	76	114
$^{50}_{27}\text{Co}$	27	23

51. $^{53}_{26}\text{Fe}^{2+}$ has 26 protons, $53 - 26 = 27$ neutrons, and two fewer electrons than protons (24 electrons) in order to have a net charge of $2+$.

52.

Symbol	Number of protons in nucleus	Number of neutrons in nucleus	Number of electrons
--------	------------------------------	-------------------------------	---------------------


CHAPTER 1 CHEMICAL FOUNDATIONS

$^{120}_{50}\text{Sn}$	50	70	50
$^{25}_{12}\text{Mg}^{2+}$	12	13	10
$^{56}_{26}\text{Fe}^{2+}$	26	30	24
$^{79}_{34}\text{Se}$	34	45	34
$^{35}_{17}\text{Cl}$	17	18	17
$^{63}_{29}\text{Cu}$	29	34	29

- 53.
- False. Neutrons have no charge; therefore, all particles in a nucleus are not charged.
 - False. The atom is best described as having a tiny dense nucleus containing most of the mass of the atom with the electrons moving about the nucleus at relatively large distances away; so much so that an atom is mostly empty space.
 - False. The mass of the nucleus makes up most of the mass of the entire atom.
 - True.
 - False. The number of protons in a neutral atom must equal the number of electrons.
54. a. True
- False; this was J. J. Thomson.
 - False; a proton is about 1800 times more massive than an electron.
 - The nucleus contains the protons and the neutrons.
55. For a $2-$ charge having 36 electrons, X^{2-} must have two fewer protons than electrons or 34 protons. This isotope of X has $79 - 34 = 45$ neutrons, and from the periodic table, X is selenium.
- True. X^{2-} has 36 electrons and 34 protons.
 - False. The isotope has 34 protons.
 - False. The isotope has 45 neutrons.
 - False. The identity is selenium, Se.
56. Because the charge of the ion is $2+$, this ion has two more protons than electrons. Therefore, the element has 88 protons, which identifies it as radium, Ra. $230 - 88 = 142$ neutrons.

CHAPTER 1 CHEMICAL FOUNDATIONS

57. Mass is conserved in a chemical reaction.



Mass: 34.0 g 12.1 g 23.3 g ?

$$\text{Mass of aluminum oxide produced} = (34.0 + 12.1) - 23.3 = 22.8 \text{ g}$$

58. Number of electrons in the unknown ion:

$$2.55 \times 10^{-26} \text{ g} \times \frac{1 \text{ kg}}{1000 \text{ g}} \times \frac{1 \text{ electron}}{9.11 \times 10^{-31} \text{ kg}} = 28 \text{ electrons}$$

Number of protons in the unknown ion:

$$5.34 \times 10^{-23} \text{ g} \times \frac{1 \text{ kg}}{1000 \text{ g}} \times \frac{1 \text{ proton}}{1.67 \times 10^{-27} \text{ kg}} = 32 \text{ protons}$$

Therefore, this ion has 32 protons and 28 electrons. This is element number 32, germanium (Ge). The net charge is 4+ because four electrons have been lost from a neutral germanium atom.

The number of electrons in the unknown atom:

$$3.92 \times 10^{-26} \text{ g} \times \frac{1 \text{ kg}}{1000 \text{ g}} \times \frac{1 \text{ electron}}{9.11 \times 10^{-31} \text{ kg}} = 43 \text{ electrons}$$

In a neutral atom, the number of protons and electrons is the same. Therefore, this is element 43, technetium (Tc).

The number of neutrons in the technetium atom:

$$9.35 \times 10^{-23} \text{ g} \times \frac{1 \text{ kg}}{1000 \text{ g}} \times \frac{1 \text{ proton}}{1.67 \times 10^{-27} \text{ kg}} = 56 \text{ neutrons}$$

The mass number is the sum of the protons and neutrons. In this atom, the mass number is 43 protons + 56 neutrons = 99. Thus this atom and its mass number is ^{99}Tc .

59. Most of the mass of the atom is due to the protons and the neutrons in the nucleus, and protons and neutrons have about the same mass ($1.67 \times 10^{-24} \text{ g}$). The ratio of the mass of the molecule to the mass of a nuclear particle will give a good approximation of the number of nuclear particles (protons and neutrons) present.


$$\frac{7.31 \times 10^{-23} \text{ g}}{1.67 \times 10^{-24} \text{ g}} = 43.8 \approx 44 \text{ nuclear particles}$$

Thus there are 44 protons and neutrons present. If the number of protons equals the number of neutrons, we have 22 protons in the molecule. One possibility would be the molecule CO_2 [$6 + 2(8) = 22$ protons].


CHAPTER 1 CHEMICAL FOUNDATIONS

Challenge Problems


60. a. One possibility is that rope B is not attached to anything and rope A and rope C are connected via a pair of pulleys and/or gears.
- b. Try to pull rope B out of the box. Measure the distance moved by C for a given movement of A. Hold either A or C firmly while pulling on the other rope.
61. a. Both compounds have C_2H_6O as the formula. Because they have the same formula, their mass percent composition will be identical. However, these are different compounds with different properties because the atoms are bonded together differently. These compounds are called isomers of each other.
- b. When wood burns, most of the solid material in wood is converted to gases, which escape. The gases produced are most likely CO_2 and H_2O .
- c. The atom is not an indivisible particle but is instead composed of other smaller particles, called electrons, neutrons, and protons.
- d. The two hydride samples contain different isotopes of either hydrogen and/or lithium. Although the compounds are composed of different isotopes, their properties are similar because different isotopes of the same element have similar properties (except, of course, their mass).
62. Because the gases are at the same temperature and pressure, the volumes are directly proportional to the number of molecules present. Let's consider hydrogen and oxygen to be monatomic gases and that water has the simplest possible formula (HO). We have the equation:




But the volume ratios are also equal to the molecule ratios, which correspond to the coefficients in the equation:




Because atoms cannot be created nor destroyed in a chemical reaction, this is not possible. To correct this, we can make oxygen a diatomic molecule:



This does not require hydrogen to be diatomic. Of course, if we know water has the formula H_2O , we get:



The only way to balance this is to make hydrogen diatomic:



63. Avogadro proposed that equal volumes of gases (at constant temperature and pressure) contain equal numbers of molecules. In terms of balanced equations, Avogadro's hypothesis (law) implies that volume ratios will be identical to molecule ratios. Assuming one molecule of octane reacting, then 1 molecule of C_xH_y produces 8 molecules of CO_2 and 9 molecules of H_2O . $C_xH_y + n O_2 \rightarrow 8 CO_2 + 9 H_2O$. Because all the carbon in octane ends up as carbon in CO_2 , octane must contain 8 atoms of C. Similarly, all hydrogen in octane ends up as hydrogen in H_2O , so one molecule of octane must contain $9 \times 2 = 18$ atoms of H. Octane formula = C_8H_{18} , and the ratio of C : H = 8 : 18 or 4 : 9.


64. From Section 1-7 of the text, the average diameter of the nucleus is about 10^{-13} cm, and the electrons move about the nucleus at an average distance of about 10^{-8} cm. From this, the diameter of an atom is about 2×10^{-8} cm.

$$\frac{2 \times 10^{-8} \text{ cm}}{1 \times 10^{-13} \text{ cm}} = 2 \times 10^5; \quad \frac{1 \text{ mi}}{1 \text{ grape}} = \frac{5280 \text{ ft}}{1 \text{ grape}} = \frac{63,360 \text{ in}}{1 \text{ grape}}$$

Because the grape needs to be 2×10^5 times smaller than a mile, the diameter of the grape would need to be $63,360 / (2 \times 10^5) \approx 0.3$ in. This is a reasonable size for a small grape.


65. Let X_a be the formula for the atom/molecule X, Y_b be the formula for the atom/molecule Y, X_cY_d be the formula of compound I between X and Y, and X_eY_f be the formula of compound

II between X and Y. Using the volume data, the following would be the balanced equations for the production of the two compounds.




From the balanced equations, $a = 2c = e$ and $b = d = 2f$.

Substituting into the balanced equations:



For simplest formulas, assume that $c = f = 1$. Thus:



Compound I = XY_2 : If X has relative mass of 1.00, $\frac{1.00}{1.00 + 2y} = 0.3043$, $y = 1.14$.

Compound II = X_2Y : If X has relative mass of 1.00, $\frac{2.00}{2.00 + y} = 0.6364$, $y = 1.14$.

The relative mass of Y is 1.14 times that of X. Thus, if X has an atomic mass of 100, then Y will have an atomic mass of 114.

CHAPTER 1 CHEMICAL FOUNDATIONS

Marathon Problem

66. a. For each set of data, divide the larger number by the smaller number to determine relative masses.

$$\frac{0.602}{0.295} = 2.04; A = 2.04 \text{ when } B = 1.00$$

$$\frac{0.401}{0.172} = 2.33; C = 2.33 \text{ when } B = 1.00$$

$$\frac{0.374}{0.320} = 1.17; C = 1.17 \text{ when } A = 1.00$$

To have whole numbers, multiply the results by 3.



Data set 1: A = 6.1 and B = 3.0

Data set 2: C = 7.0 and B = 3.0

Data set 3: C = 3.5 and A = 3.0 or C = 7.0 and A = 6.0

Assuming 6.0 for the relative mass of A, the relative masses would be A = 6.0, B = 3.0, and C = 7.0 (if simplest formulas are assumed).

- b. Gas volumes are proportional to the number of molecules present. There are many possible correct answers for the balanced equations. One such solution that fits the gas volume data is:



In any correct set of reactions, the calculated mass data must match the mass data given initially in the problem. Here, the new table of relative masses would be:

$$\frac{6(\text{mass } A_2)}{\text{mass } B_4} = \frac{0.602}{0.295}; \text{mass } A_2 = 0.340(\text{mass } B_4)$$

$$\frac{4(\text{mass } C_3)}{\text{mass } B_4} = \frac{0.401}{0.172}; \text{mass } C_3 = 0.583(\text{mass } B_4)$$

$$\frac{2(\text{mass } C_3)}{3(\text{mass } A_2)} = \frac{0.374}{0.320}; \text{mass } A_2 = 0.570(\text{mass } C_3)$$

CHAPTER 1 CHEMICAL FOUNDATIONS

Assume some relative mass number for any of the masses. We will assume that mass B = 3.0, so mass B₄ = 4(3.0) = 12.

Mass C₃ = 0.583(12) = 7.0, mass C = 7.0/3

Mass A₂ = 0.570(7.0) = 4.0, mass A = 4.0/2 = 2.0

When we assume a relative mass for B = 3.0, then A = 2.0 and C = 7.0/3. The relative masses having all whole numbers would be A = 6.0, B = 9.0, and C = 7.0.

Note that any set of balanced reactions that confirms the initial mass data is correct. This is just one possibility.